

AK

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/00	A1	(11) International Publication Number: WO 98/32421 (43) International Publication Date: 30 July 1998 (30.07.98)
(21) International Application Number: PCT/US98/01649 (22) International Filing Date: 29 January 1998 (29.01.98) (30) Priority Data: 08/794,690 29 January 1997 (29.01.97) US (71) Applicant: BAUSCH & LOMB INCORPORATED (US/US); One Bausch & Lomb Place, Rochester, NY 14604-2701 (US). (72) Inventors: HU, Zhenze; 28 Wenham Lane, Pittsford, NY 14534 (US). DENICK, John; 290 Tobey Road, Pittsford, NY 14534 (US). (74) Agents: KONKOL, Chris, P. et al.; Bausch & Lomb Incorporated, One Bausch & Lomb Place, Rochester, NY 14604-2701 (US).	(81) Designated States: AL, AU, BB, BR, BY, CA, CU, EE, ES, FI, GB, HU, IL, JP, KE, KG, KP, KR, MD, MK, MN, MW, NO, NZ, PL, PT, RU, SE, SI, SK, TJ, TR, TT, UA, UZ, VN, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: OPHTHALMIC COMPOSITIONS INCLUDING GLYCERIN AND PROPYLENE GLYCOL (57) Abstract <p>There are disclosed ophthalmic compositions having high water-binding properties which are useful as: moisturizing and lubricating (i.e. artificial tear) solutions, dry eye therapies, contact lens wetting and lubricating solutions, and as delivery vehicles for ophthalmic drugs. The subject compositions include glycerin in combination with propylene glycol. The subject compositions may further include cellulose derivatives, e.g. hydroxypropyl methyl cellulose, along with preservatives, e.g. benzylalkonium chloride, PHMB, sorbic acid, etc.. Preferred compositions have at least 11 % bound water, a pH from about 7.1 to 7.5, and an osmolality between about 280 to about 320 mOsm/Kg.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UC	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CJ	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

OPHTHALMIC COMPOSITIONS INCLUDING GLYCERIN AND PROPYLENE GLYCOL

FIELD OF THE INVENTION

The present invention is directed toward ophthalmic compositions, particularly those provided as buffered, aqueous solutions. The subject compositions are useful as: moisturizing and lubricating eye drops, delivery vehicles for ophthalmic drugs, and as contact lens wetting and lubricating solutions.

BACKGROUND

Ophthalmic compositions used for the treatment of "dry eye" symptoms include demulcents (or humectants) for lubricating mucous membrane surfaces and for relieving dryness and irritation. The term "demulcent", as used herein is intended to mean an agent, usually a water-soluble polymer, which is applied topically to the eye to protect and lubricate mucous membrane surfaces and relieve dryness and irritation. Within this meaning, the terms "humectant" and "wetting agent" are also commonly used. Furthermore, it will be understood that some constituents possess several functional attributes. For example, cellulose derivatives are common demulcents, but are also used as "viscosity increasing agents". Similarly, glycerin is a known demulcent but is also used as a "tonicity adjusting agent". Examples of the most widely used demulcents include: polyvinyl alcohol, polyvinyl pyrrolidone, cellulose derivatives and polyethylene glycol.

"Over-the-counter" use of demulcents within ophthalmic compositions is regulated by the US Food & Drug Administration. For example, the Federal Register (21 CFR Part 349) entitled Ophthalmic Drug Products for Over-the-Counter Use: Final Monograph list the accepted demulcents along with appropriate concentration ranges for each. Specifically, § 349.12 list the following approved "monograph" demulcents: (a) cellulose derivatives: (1) carboxymethylcellulose sodium, (2) hydroxyethyl cellulose, (3) hydroxypropyl methylcellulose, methylcellulose, (b) dextran 70, (c) gelatin, (d) polyols, liquid: (1) glycerin, (2) polyethylene glycol 300, (3) polyethylene glycol 400, (4) polysorbate 80, (5) propylene glycol (e) polyvinyl alcohol, and (f) povidone (polyvinyl

pyrrolidone). § 349.30 further provides that in order to fall within the monograph, no more than three of the above-identified demulcents may be combined.

Specific examples of known ophthalmic compositions including various demulcents are provided below.

US Patent No. 5,591,426 to Dabrowski et al. discloses a ophthalmic solution useful as an artificial tear. The reference includes a specific example of a borate buffered, preserved (e.g. benzalkonium chloride), aqueous solution including the following three demulcents: 1) glycerin, 2) polyvinyl pyrrolidone, and 3) a cellulose derivative, e.g. hydroxypropyl methyl cellulose.

US Patent No. 5,106,615 to Dikstein discloses isotonic humectant eyedrops including glycerin, polyethylene glycol, or propylene glycol with an anionic polymer such as Carbomer 941.

US Patent No. 2,703,777 to Feinstein et al. generally describes a preserved, buffered, isotonic ophthalmic gel including: 1) a humectant, preferably glycerin (sorbitol and propylene glycol are also listed); 2) methyl cellulose, and 3) polyethylene glycol.

US Patent No. 4,029,817 to Blanco et al. discloses a contact lens preserving solution including propylene glycol in combination with polysorbate 80 and/or polyvinyl pyrrolidone. Similarly, US Patent No. 5,141,665 to Sherman discloses a contact lens cleaning, wetting and storing solution which includes propylene glycol as a wetting agent. Also, US Patent No. 4,525,346 to Stark discloses a borate buffered, preserved contact lens solution including propylene glycol.

US Patent Nos. 3,767,788; 3,767,789; 3,856,919; 3,907,985; 3,920,810; 3,947,573; 3,987,163 all to Billy Rankin disclose ophthalmic solutions for the treatment of "dye eye". These references generally teach the use of polyethylene oxide, polystyrene sulfonate, and polyacrylamide, with polyalkylene glycols, e.g. polyethylene glycol or

propylene glycol. These references include specific example solutions including several demulcents combined with one another, namely, 1) polyethylene glycol, 2) polyvinyl pyrrolidone and a 3) cellulose derivative, e.g. hydroxy ethyl cellulose.

US Patent No. 3,549,747 to Krezanoski et al. discloses a preserved contact lens wetting solution including polyvinyl alcohol with a cellulose derivative, e.g. hydroxy ethyl cellulose. Similarly, US Patent No. 4,131,651 to Shah et al. discloses an ophthalmic solution for the treatment of dry eye which includes polyvinyl alcohol with a cellulose derivative. US Patent No. 4,120,949 to Bapatla et al. discloses a preserved ophthalmic solution including 1) polyvinyl alcohol, 2) polyvinylpyrrolidone, and 3) one or more cellulose derivatives. Also similarly, US Patent No. 4,409,205 to Shively discloses a specific example of a preserved ophthalmic solution including: polyvinyl alcohol, polyethylene glycol 6000, and dextrose. This reference also generally discloses the use of tonicity adjusting agents selected from the group of: mannitol, sorbitol, dextrose, sucrose, urea, and glycerin.

New ophthalmic compositions are sought which provide greater relief from dryness and irritation. In this regard, compositions capable of binding higher amounts of water have been identified by applicants as being particularly preferred. Furthermore, it is desired to accomplish these improvements while using approved monograph demulcents.

SUMMARY OF THE INVENTION

The present invention is an ophthalmic composition comprising glycerin and propylene glycol. It has been discovered that propylene glycol and glycerin are capable of binding significantly greater amounts of water than other demulcents, as will be described below.

In one preferred embodiment, the subject composition is provided as a buffered, aqueous solution which includes a third demulcent, preferably a cellulose derivative. The subject composition may be unpreserved (provided in a single dose format), or may be preserved, e.g. with benzylalkonium chloride, PHMB, sorbic acid, etc.

DETAILED DESCRIPTION OF THE INVENTION

As previously described, the subject composition finds particular utility as a moisturizing and lubricating eye drop (i.e. an artificial tear solution), a delivery vehicle for ophthalmic drugs, and as a contact lens wetting and lubricating solution. In most of these applications, the subject composition is provided as a buffered aqueous solution. Such a solution typically has a viscosity from about 1 to about 50 cps. As a solution, the subject composition is usually dispensed in the eye in the form of an eye drop. It should be understood, however, that the subject composition may also be formulated as a viscous liquid (i.e. viscosities from 50 to several thousand cps), gel, or ointment. Furthermore, in some contact lens related embodiments, lenses may be soaked or otherwise exposed to the subject composition prior to wear.

The present ophthalmic compositions include glycerin and propylene glycol. In addition to these two demulcents, other demulcents may also be used. However, the FDA regulations limit the number of demulcents which may be used together in over-the-counter ophthalmics, to three demulcents. In light of this regulation, and in an effort to simplify formulations and reduce undesirable interaction among constituents, it may be desirable to limit the number of demulcents used in a given composition. Thus, in several preferred embodiments of the subject invention, the only demulcents used are glycerin and propylene glycol, or alternatively glycerin, propylene glycol and one other demulcent, preferably a cellulose derivative. Cellulose derivatives are commonly used to increase viscosity, and as such, offer other advantages. Specific cellulose derivatives include: hydroxypropyl methyl cellulose, carboxymethyl cellulose, methyl cellulose, hydroxyethyl cellulose, etc. In these embodiments, other demulcents such as polyvinyl pyrrolidone, polyvinyl alcohol, polyethylene glycol, and other components such as polyethylene oxide, and polyacrylic acid are specifically excluded.

In still other embodiments, other or additional demulcents may be used in combination with glycerin and propylene glycol. For example, polyvinyl pyrrolidone, polyvinyl alcohol, may also be used.

The demulcents used in the present invention are used in effective amounts (i.e. "demulcifying amounts") for providing a demulcifying effect, i.e. sufficient to lubricating mucous membrane surfaces and to relieve dryness and irritation. The specific quantities of demulcents used in the present invention will vary depending upon the application. However, typically the amount of glycerin is suitably 0.05 to 1.5 weight percent (%w/w), preferably from about 0.2 to about 1.0%; typically the amount of propylene glycol is from about 0.2 to about 1.5%, but preferably about 1% (w/w). In one embodiment of the invention, the ratio (w/w) of propylene glycol to glycerin in the composition is 2:1 to 10:1, preferably 3:1 to 5:1 (w/w). A composition containing this ratio of propylene glycol to glycerin has been found to provide improved comfort.

In one embodiment of the invention, a cellulose derivative may be included in the amount of from about 0.2 to about 3%, but preferably about 0.5% (w/w). A preferred cellulose derivative is pharmaceutical grade hydroxypropyl methylcellulose (HPMC), such as Methocel E 15 LV - premium, available from Dow Chemical Company.

When used, any pharmaceutically acceptable buffer system may be utilized; however, a preferred buffer system is provided by sodium borate and boric acid in amounts necessary to produce a pH of about 6.0 to about 8.0, but more preferably from about 7.1 to about 7.5.

The composition may be designed for a variety of osmolalities, but in most applications, iso-osmolal (with respect to the fluids of the eye) compositions are preferred. Osmolalities typically range from about 175 to about 350 mOsm/kg, preferably 250 to 330 mOsm/kg, but more preferably from about 280 to about 320 mOsm/kg. The osmolality of the solution may be adjusted by means of well known osmolality adjusting agents, e.g. sodium chloride and potassium chloride, and monosaccharides.

As previously indicated, the subject composition may include a preservative in an amount effective to preserve the solution. As is known in the art, the amount of

preservative required will vary upon the specific preservative and the application, e.g. moisturizing eye drop, contact lens solution, etc. For non-contact lens applications, benzalkonium chloride (BAK) is a preferred preservative typically used in concentrations from about 0.01 to about 0.10 % (w/w). BAK is a well known preservative which comprises a mixture of alkyldimethyl benzylammonium chlorides. For contact lens applications, other preservative are more preferred, such as sorbic acid, PHMB, and other polyquats. Alternatively, the subject compositions may be preservative-free.

The composition may include a number of additional components. For example, the solution may include edetate disodium as a co-preservative and/or chelating agent.

EXAMPLE I

As an illustration of the present invention, a preferred moisturizing eye drop formulation is provided below.

TABLE I

Constituent	%w/w
glycerin	1.0
propylene glycol	0.5
hydroxypropylmethyl cellulose (HPMC)	1.0
boric acid	0.300
sodium borate	0.035
sodium chloride (NaCl)	0.096
potassium chloride (KCl)	0.097
edetate disodium (EDTA)	0.030
benzalkonium chloride (BAK) (50%)	0.021
purified water	q.s. to 100%

The formulation was prepared by adding each of NaCl, KCl, sodium borate, boric acid, EDTA and HPMC (Type E15-LV Premium) sequentially to a volume of heated water (80-90°C) that amounted to 70-85% of the final batch volume. This addition was accomplished under constant agitation and each component was allowed to dissolve or disperse before adding the next. The resulting solution was mixed for about 30 minutes and maintained at a temperature of about 85°C. With continued agitation, the batch was cooled to 50°C (+/-5°C), at which point propylene glycol, glycerin, and BAK (50% solution) were each sequentially added to the solution. The batch was cooled under

agitation to about 20°C (+/-5°C). The pH of the solution was then adjusted to about 7.1 to about 7.5 using increments of either 1N NaOH or 1N HCl, and was then brought to final volume with 20-30°C water and mixed for at least 15 minutes. The BAK concentration is finally adjusted to a 95-110 ppm range.

EXAMPLE II

As a further illustration of the present invention, another preferred moisturizing eye drop formulation is provided below.

TABLE II

Constituent	%w/w
glycerin	0.300
propylene glycol	1.000
boric acid	0.300
sodium borate	0.035
sodium chloride (NaCl)	0.129
potassium chloride (KCl)	0.125
edetate disodium (EDTA)	0.030
benzalkonium chloride (BAK) (50%)	0.021
purified water	q.s. to 100%

The formulation may be prepared as follows. Each of NaCl, KCl, sodium borate, boric acid, and EDTA is sequentially added to a volume of heated water (80-90°C) that amounts to 70-85% of the final batch volume. This addition is accomplished under constant agitation, and each component is allowed to dissolve or disperse before adding the next. The resulting solution is mixed for about 30 minutes and maintained at a temperature of about 85°C. With continued agitation, the batch is allowed to cool to 50°C (+/-5°C), at which point propylene glycol, glycerin, and BAK (50% solution) is each sequentially added to the solution. The batch is then cooled under agitation to about 20°C (+/-5°C), the pH is adjusted to about 7.1 to about 7.5 using increments of either 1N NaOH or 1N HCl, and the final volume is obtained with 20-30°C water, mixing for at least 15 minutes. The BAK concentration is finally adjusted to a 95-110 ppm range.

EXAMPLE III

The water-binding capacity of several demulcents were evaluated based upon the freezably water fraction of a 10% demulcent solution. Samples were evaluated using differential scanning calorimetry (DSC), utilizing a TA instrument DSC 912 with a 2100 controller. The instrument was calibrated using indium metal and HPLC grade water for the upper and lower temperatures corrections, respectively.

The indium metal calibration was completed as follows. An aluminum sample pan and cover were weighed. Once the pan weight was zeroed, a piece of indium metal was placed in the pan and covered and weighed. Approximately 10 mg of indium was used for both cells. The aluminum pan was hermetically sealed and placed on the appropriate cell head, A or B. An empty aluminum pan was hermetically sealed and placed on the reference head. Nitrogen gas was purged through the cell entering through the vacuum and cooling ports at a fixed rate (50 on the scale of a Gilmont flowmeter). The cell was closed and the following temperature program was used:

1. equilibrate at 130°C;
2. isothermal for 4.0 minutes;
3. ramp 1.00°C/minute to 180°C;
4. isothermal for 2.00 minutes.

The HPLC water calibration was completed as follows. Two water samples were prepared in a fashion similar to that as described above. Approximately 10µl of HPLC grade water was injected into a zeroed pan assembly. The weight was obtained with the cover on the pan so that the sample did not evaporate while being weighed. The following temperature program was used for all water samples.

1. Equilibrate at -20°C;
2. ramp 1.00°C/minutes to 20.0°C.

Indium endotherms were integrated between 156.3°C and 158.3 °C. Water endotherms were integrated from -1°C to 5°C. The onset of the melt endotherms were averaged from cell A and B. The average melt temperature for the indium and water calibration samples were imputed into the temperature calibration. Cell constants were calculated for Cell A and B using the delta H values determined for the water samples. The literature value for the heat of fusion of water is 333.6J/g. Cell constants were calculated

by dividing this value by the delta H value obtained on that cell. These values were then inputted into the module parameters.

The demulcent samples were tested in a similar manner as the water calibration samples described above. Three runs for each sample were completed. The demulcent sample endotherms were integrated between approximately -15°C and 7°C. The results of the evaluation are provided in Table III, below.

TABLE III

DEMULCENT	grams bound water per gram of dry demulcent
methylcellulose	1.237
dextran 70	0.683
hydroxypropyl methyl cellulose	1.138
polyvinylpyrrolidone	1.370
carboxymethylcellulose sodium	0.947
hydroxyethylcellulose	1.168
polyvinyl alcohol	0.956
polysorbate 80	0.982
polyethylene glycol 300	1.959
polyethylene glycol 400	1.907
glycerin	2.212
gelatin	1.340
propylene glycol	2.756

As indicated from the data of Table III, propylene glycol and glycerin had far greater water-binding capacities than all other demulcent tested. This result was unexpected, particularly given the infrequent use of propylene glycol as a demulcent in the art. By combining glycerin and propylene glycol together as demulcents, it is possible to prepare improved ophthalmic compositions having at least 11%, and preferably about 13% percent bound water as defined by Formula I below. This has been accomplished while complying with the monograph requirements.

FORMULA I

$$\text{Percent Bound Water} = \text{Percent Total Water} - 100 (\Delta H_{\text{sample}} / \Delta H_{\text{water}})$$

CLAIMS

1. An ophthalmic composition comprising an effective demulcifying amount of:
(a) glycerin, and (b) propylene glycol.
2. The composition of claim 1 including at least one of : a cellulose derivative, dextran, gelatin, polyethylene glycol, polysorbate, polyvinyl alcohol and polyvinylpyrrolidone.
3. The composition of claim 2 including at least one cellulose derivative selected from the group: hydroxypropyl methyl cellulose, carboxymethyl cellulose, methyl cellulose, and hydroxyethyl cellulose.
4. The composition of claim 1 including a preservative.
5. The composition of claim 4 including at least one preservative selected from the group: benzylalkonium chloride, polyquat, PHMB, and sorbic acid.
6. The composition of claim 1 having a pH from about 7.1 to about 7.5, an osmolality between about 280 to about 320 mOsm/Kg, and at least about 11 percent bound water.
7. An ophthalmic solution useful as an artificial tear comprising:
 - (a) from about 0.2 to about 1.5 weight percent of glycerin,
 - (b) from about 0.2 to about 1.5 weight percent of propylene glycol,
 - (c) from about 0.2 to 3.0% of a cellulose derivative, and
 - (d) a borate buffer, andwherein said solution has an osmolality between about 280 to about 320 mOsm/Kg, and a pH from about 7.1 to about 7.5.

8. The solution of claim 7 which excludes polyvinylpyrrolidone, polyvinyl alcohol, polyethylene glycol, and polyethylene oxide, polyacrylic acid, anionic polymers.

9. A method for moisturizing or lubricating the eye by contacting the eye with an ophthalmic composition comprising an effective demulcifying amount of: (a) glycerin and (b) propylene glycol.

10. A method for wetting a contact lens by contacting the eye with an ophthalmic composition comprising an effective demulcifying amount of: (a) glycerin and (b) propylene glycol.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/01649

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K9/00

According to International Patent Classification(IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 92 00707 A (INSITE VISION INCORPORATED) 23 January 1992 see claims 1-3,9-11 see page 15, line 24 - line 30 see page 18, line 4 - line 23 see page 21, line 20 - page 22, line 2	1-10
X	EP 0 580 074 A (ASTA MEDICAKTIENGESELLSCHAFT) 26 January 1994 see page 5; example 7 -/-	1,4,5,9, 10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"A" document member of the same patent family

Date of the actual completion of the international search

11 May 1998

Date of mailing of the international search report

20/05/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Ventura Amat, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/01649

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p> DATABASE WPI Section Ch, Week 9502 Derwent Publications Ltd., London, GB; Class A96, AN 95-011747 XP002064494 & JP 06 298 649 A (TEIKA SEIYAKU) , 25 October 1994 see abstract </p>	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/01649

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9200707 A	23-01-1992	US 5188826 A	23-02-1993
		AU 8184991 A	04-02-1992
		CA 2086120 A	07-01-1992
		EP 0538313 A	28-04-1993
EP 580074 A	26-01-1994	JP 6040949 A	15-02-1994
		AT 117207 T	15-02-1995
		DE 69300054 D	02-03-1995
		DE 69300054 T	30-05-1996
		ES 2068725 T	16-04-1995